

- FLEISCHMANNIA ANTIQUORUM (Standl. & Steyerem.) R. M. King & H. Robinson, comb. nov. *Eupatorium antiquorum* Standl. & Steyerem., Publ. Field Mus. Nat. Hist. Chic. Bot. 22: 302. 1940.
- FLEISCHMANNIA PASTAZAE (B.L.Robinson) R. M. King & H. Robinson, comb. nov. *Eupatorium pastazae* B. L. Robinson, Biblioth. Bot. 29: 160 Heft. 116. 1937.
- FLEISCHMANNIA SAXORUM (Standl. & Steyerem.) R. M. King & H. Robinson, comb. nov. *Eupatorium saxorum* Standl. & Steyerem., Publ. Field Mus. Nat. Hist. Chic. Bot. 23: 189. 1944.
- GRAZIELIA MOLLICOMA (B.L.Robinson) R. M. King & H. Robinson, comb. nov. *Eupatorium mollicomum* B. L. Robinson, Contrib. Gray Herb. n.s. 68: 26. 1923.
- GUAYANIA CRASSICAULIS (Steyerem.) R. M. King & H. Robinson, comb. nov. *Eupatorium crassicaule* Steyerem., Fieldiana, Bot. 28: 629, No. 3. 1953.
- HEBECLINIUM OBTUSISQUAMOSUM (Hieron. ex Sod.) R. M. King & H. Robinson, comb. nov. *Eupatorium obtusisquamosum* Hieron. ex Sod., Bot. Jahrb. 29: 14. 1900.
- HETEROCONDYLUS ALATUS (Vellozo) R. M. King & H. Robinson, comb. nov. *Chrysocoma alata* Vellozo, Fl. Flum. 313. 1825. The Vellozo name has priority for the species commonly known as *Eupatorium vautherianum* DC.
- KOANOPHYLLON GRANDICEPS (Wright) R. M. King & H. Robinson, comb. nov. *Eupatorium grandiceps* Wright, Anales Acad. Ci. Med. Habana 6: 178. 1869. Sauv. Fl. Cub. 76. 1873.
- KOANOPHYLLON GRISEBACHIANUM (Alain) R. M. King & H. Robinson, comb. nov. *Eupatorium grisebachianum* Alain, Candollea 17: 121. 1960. The name was provided by Alain for *Eupatorium incisum* Griseb. not A.Rich.
- KOANOPHYLLON QUISQUEYANUM (Alain) R. M. King & H. Robinson, comb. nov. *Eupatorium quisqueyanum* Alain, Moscosoa 1 (1): 48. 1976.
- NEOCABRERIA PENNIVENIA (B.L.Robinson) R. M. King & H. Robinson, comb. nov. *Symphopappus pennivenius* B. L. Robinson, Contrib. Gray Herb. n.s. 68: 7. 1923.
- PHALACRAEA LONGIPETIOLATA (B.L.Robinson) R. M. King & H. Robinson, comb. nov. *Piqueria longipetiolata* B. L. Robinson, Proc. Amer. Acad. 43: 27. 1907.

- SPHAEREUPATORIUM SPHAEROCEPHALUM (Sch.Bip. ex Baker) R. M. King
& H. Robinson, comb. nov. *Eupatorium sphaerocephalum* Sch.
Bip. ex Baker in Martius, Fl. Bras. 6 (2): 317. 1876.
- STEVIOPSIS VIGINTISETA (DC.) R. M. King & H. Robinson, comb. nov.
Stevia vigintisetata DC., Prodr. 5: 123. 1836.
- One species transferred to the tribe Inuleae also needs a new
combination:
- IPHIONA RETOFRACTA (Thunb.) H. Robinson, comb. nov. *Eupatorium*
retrofractum Thunb., Prodr. Pl. Cap. 142. 1800.

STUDIES IN THE EUPATORIEAE (ASTERACEAE). CCVIII.

ADDITIONS TO *BADILLOA* AND *BARTLETTINA* FROM ECUADOR

R. M. King and H. Robinson
Department of Botany

Smithsonian Institution, Washington, D.C., 20560.

Material recently sent on loan by the New York Botanical Garden contains specimens of one previously described species needing a new combination and one species that is undescribed.

BADILLOA ATRESCENS (B.L.Robinson) R. M. King & H. Robinson, comb. nov. *Eupatorium atrescens* B.L.Robinson, Contrib. Gray Herb. n.s. 77: 9. 1926. The species was originally described as a shrub, 8 ft. tall with purple flowers between San Lucas and Oña in the Province of Loja, Ecuador. When examined many years ago, the species seemed to represent an intergeneric hybrid between *Badilloa* and *Aristeguietia* with the essential key characters of the former. We deferred making a combination without evidence of the existence of an established population. The following specimen provides that evidence. ECUADOR: Azuay-"Oriente" Border: Eastern Cordillera, between Oña and the río Yacuambi. West slope, 8,000-9,500 ft. elev. Vine. Leaves deep green above, pale below. Bracts green suffused with reddish purple. Base of corolla cream; upper part of tube, lobes and style branches dull lavender-purple. Sept. 10-19, 1945. F. Prieto P-213.

BARTLETTINA CAMPII R.M.King & H.Robinson, sp. nov.

Plantae scandentes 5 m longae mediocriter ramosae. Caules pallide brunnescentes subhexagonales dense grosse hirsuti. Folia opposita, petiolis 1.5-4.5 cm longis; laminae triangulares plerumque 4-10 cm longae et 3-7 cm latae base valde cordatae margine multo crenato-dentatae apice breviter anguste acuminatae supra ubiquiter pilosae subtus plerumque in nervis prominule reticulatis pilosae ad basem distincte trinervatae. Inflorescentiae late laxae pyramidaliter paniculatae in ramis distalis congestae, ramis ultimis 2-11 mm longis hirsutis. Capitula late campanulata 8-10 mm alta et 6-10 mm lata; squamae involucri ca. 50 distincte subimbricatae valde inaequilongae ca. 4-seriatae ovatae vel lineari-lanceolatae 2-6 mm longae et 0.3-1.3 mm latae late vel anguste acutae margine puberulo-fimbriatae extus puberulae vel glabrae 2-4-costatae; receptacula lenitèr convexa sparse pilosula. Flores ca. 40 in capitulo; corollae purpureae ca. 5.0-5.5 mm longae extus glabrae, tubis cylindraceis ca. 3 mm longis, faucibus infundibularibus ca. 2 mm longis, lobis triangularibus ca. 0.5 mm longis et base 0.4-0.5 mm latis utrinque laevibus; filamenta in parte superiore ca. 0.5 mm longa, cellulis plerumque